

## Section 1.2: Row Reduction and Echelon Forms

### Echelon form (or row echelon form):

1. All nonzero rows are above any rows of all zeros.
2. Each *leading entry* (i.e. left most nonzero entry) of a row is in a column to the right of the leading entry of the row above it.
3. All entries in a column below a leading entry are zero.

### EXAMPLE: Echelon forms

$$\begin{array}{l}
 \text{(a)} \left[ \begin{array}{ccccc}
 \blacksquare & * & * & * & * \\
 0 & \blacksquare & * & * & * \\
 0 & 0 & 0 & 0 & 0 \\
 0 & 0 & 0 & 0 & 0
 \end{array} \right]
 \end{array}
 \quad
 \begin{array}{l}
 \text{(b)} \left[ \begin{array}{ccc}
 \blacksquare & * & * \\
 0 & \blacksquare & * \\
 0 & 0 & \blacksquare \\
 0 & 0 & 0
 \end{array} \right]
 \end{array}$$

$$\text{(c)} \left[ \begin{array}{cccccccccccc}
 0 & \blacksquare & * & * & * & * & * & * & * & * & * & * \\
 0 & 0 & 0 & \blacksquare & * & * & * & * & * & * & * & * \\
 0 & 0 & 0 & 0 & \blacksquare & * & * & * & * & * & * & * \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & \blacksquare & * & * & * & * \\
 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \blacksquare & * & * & *
 \end{array} \right]$$

**Reduced echelon form:** Add the following conditions to conditions 1, 2, and 3 above:

4. The leading entry in each nonzero row is 1.
5. Each leading 1 is the only nonzero entry in its column.

**EXAMPLE** (continued):

Reduced echelon form :

$$\begin{bmatrix} 0 & 1 & * & 0 & 0 & * & * & 0 & 0 & * & * \\ 0 & 0 & 0 & 1 & 0 & * & * & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & 1 & * & * & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & * & * \end{bmatrix}$$

**Theorem 1 (Uniqueness of The Reduced Echelon Form):**

Each matrix is row-equivalent to one and only one reduced echelon matrix.

## Important Terms:

- **pivot position:** a position of a leading entry in an echelon form of the matrix.
- **pivot:** a nonzero number that either is used in a pivot position to create 0's or is changed into a leading 1, which in turn is used to create 0's.
- **pivot column:** a column that contains a pivot position.

(See the Glossary at the back of the textbook.)

**EXAMPLE:** Row reduce to echelon form and locate the pivot columns.

$$\begin{bmatrix} 0 & -3 & -6 & 4 & 9 \\ -1 & -2 & -1 & 3 & 1 \\ -2 & -3 & 0 & 3 & -1 \\ 1 & 4 & 5 & -9 & -7 \end{bmatrix}$$

**Solution**

pivot

$$\begin{bmatrix} 1 & 4 & 5 & -9 & -7 \\ -1 & -2 & -1 & 3 & 1 \\ -2 & -3 & 0 & 3 & -1 \\ 0 & -3 & -6 & 4 & 9 \end{bmatrix}$$

↑  
pivot column

$$\begin{bmatrix} 1 & 4 & 5 & -9 & -7 \\ 0 & 2 & 4 & -6 & -6 \\ 0 & 5 & 10 & -15 & -15 \\ 0 & -3 & -6 & 4 & 9 \end{bmatrix}$$

Possible Pivots:

$$\begin{bmatrix} 1 & 4 & 5 & -9 & -7 \\ 0 & 2 & 4 & -6 & -6 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -5 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 4 & 5 & -9 & -7 \\ 0 & 2 & 4 & -6 & -6 \\ 0 & 0 & 0 & -5 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Original Matrix:

$$\begin{bmatrix} 0 & -3 & -6 & 4 & 9 \\ -1 & -2 & -1 & 3 & 1 \\ -2 & -3 & 0 & 3 & -1 \\ 1 & 4 & 5 & -9 & -7 \end{bmatrix}$$

$\uparrow \quad \uparrow \quad \uparrow$   
 pivot columns:    1   2        4

**Note:** There is no more than one pivot in any row. There is no more than one pivot in any column.

**EXAMPLE:** Row reduce to echelon form and then to reduced echelon form:

$$\begin{bmatrix} 0 & 3 & -6 & 6 & 4 & -5 \\ 3 & -7 & 8 & -5 & 8 & 9 \\ 3 & -9 & 12 & -9 & 6 & 15 \end{bmatrix}$$

**Solution:**

$$\begin{bmatrix} 0 & 3 & -6 & 6 & 4 & -5 \\ 3 & -7 & 8 & -5 & 8 & 9 \\ 3 & -9 & 12 & -9 & 6 & 15 \end{bmatrix} \sim \begin{bmatrix} 3 & -9 & 12 & -9 & 6 & 15 \\ 3 & -7 & 8 & -5 & 8 & 9 \\ 0 & 3 & -6 & 6 & 4 & -5 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & -9 & 12 & -9 & 6 & 15 \\ 0 & 2 & -4 & 4 & 2 & -6 \\ 0 & 3 & -6 & 6 & 4 & -5 \end{bmatrix}$$

Cover the top row and look at the remaining two rows for the left-most nonzero column.

$$\begin{bmatrix} 3 & -9 & 12 & -9 & 6 & 15 \\ 0 & 2 & -4 & 4 & 2 & -6 \\ 0 & 3 & -6 & 6 & 4 & -5 \end{bmatrix} \sim \begin{bmatrix} 3 & -9 & 12 & -9 & 6 & 15 \\ 0 & 1 & -2 & 2 & 1 & -3 \\ 0 & 3 & -6 & 6 & 4 & -5 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & -9 & 12 & -9 & 6 & 15 \\ 0 & 1 & -2 & 2 & 1 & -3 \\ 0 & 0 & 0 & 0 & 1 & 4 \end{bmatrix} \text{ (echelon form)}$$

### Final step to create the reduced echelon form:

Beginning with the rightmost leading entry, and working upwards to the left, create zeros above each leading entry and scale rows to transform each leading entry into 1.

$$\begin{bmatrix} 3 & -9 & 12 & -9 & 0 & -9 \\ 0 & 1 & -2 & 2 & 0 & -7 \\ 0 & 0 & 0 & 0 & 1 & 4 \end{bmatrix} \sim \begin{bmatrix} 3 & 0 & -6 & 9 & 0 & -72 \\ 0 & 1 & -2 & 2 & 0 & -7 \\ 0 & 0 & 0 & 0 & 1 & 4 \end{bmatrix}$$
$$\sim \begin{bmatrix} 1 & 0 & -2 & 3 & 0 & -24 \\ 0 & 1 & -2 & 2 & 0 & -7 \\ 0 & 0 & 0 & 0 & 1 & 4 \end{bmatrix}$$

## SOLUTIONS OF LINEAR SYSTEMS

- **basic variable:** any variable that corresponds to a pivot column in the augmented matrix of a system.
- **free variable:** all nonbasic variables.

### EXAMPLE:

$$\begin{bmatrix} 1 & 6 & 0 & 3 & 0 & 0 \\ 0 & 0 & 1 & -8 & 0 & 5 \\ 0 & 0 & 0 & 0 & 1 & 7 \end{bmatrix} \quad \begin{array}{rcl} x_1 + 6x_2 & + 3x_4 & = 0 \\ & x_3 - 8x_4 & = 5 \\ & & x_5 = 7 \end{array}$$

pivot columns:  $1, 3, 5$

basic variables:  $x_1, x_3, x_5$

free variables:  $x_2, x_4$



**Final Step in Solving a Consistent Linear System:** After the augmented matrix is in **reduced** echelon form and the system is written down as a set of equations:

*Solve each equation for the basic variable in terms of the free variables (if any) in the equation.*

**EXAMPLE:**

$$\begin{array}{rcccc} x_1 & +6x_2 & & +3x_4 & = & 0 \\ & & x_3 & -8x_4 & = & 5 \\ & & & & x_5 & = & 7 \end{array} \quad \left\{ \begin{array}{l} x_1 = -6x_2 - 3x_4 \\ x_2 \text{ is free} \\ x_3 = 5 + 8x_4 \\ x_4 \text{ is free} \\ x_5 = 7 \end{array} \right. \quad \text{(general solution)}$$

The **general solution** of the system provides a parametric description of the solution set. (The free variables act as parameters.) The above system has **infinitely many solutions**.

Why? *Because it has free variables, which can take on any values.*

**Warning: Use only the reduced echelon form to solve a system.**

## Existence and Uniqueness Questions

### EXAMPLE:

$$\begin{bmatrix} & 3x_2 & -6x_3 & +6x_4 & +4x_5 & = & -5 \\ 3x_1 & -7x_2 & +8x_3 & -5x_4 & +8x_5 & = & 9 \\ 3x_1 & -9x_2 & +12x_3 & -9x_4 & +6x_5 & = & 15 \end{bmatrix}$$

In an earlier example, we obtained the echelon form:

$$\begin{bmatrix} 3 & -9 & 12 & -9 & 6 & 15 \\ 0 & 2 & -4 & 4 & 2 & -6 \\ 0 & 0 & 0 & 0 & 1 & 4 \end{bmatrix} \quad (x_5 = 4)$$

No equation of the form  $0 = c$ , where  $c \neq 0$ , so the system is consistent.

**Free variables:**  $x_3$  and  $x_4$

**Consistent system  
with free variables**

$\Rightarrow$  infinitely many solutions.

**EXAMPLE:**

$$\begin{array}{rcl} 3x_1 + 4x_2 & = & -3 \\ 2x_1 + 5x_2 & = & 5 \\ -2x_1 - 3x_2 & = & 1 \end{array} \rightarrow \begin{bmatrix} 3 & 4 & -3 \\ 2 & 5 & 5 \\ -2 & -3 & 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 3 & 4 & -3 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow \begin{array}{l} 3x_1 + 4x_2 = -3 \\ x_2 = 3 \end{array}$$

**Consistent system,  
no free variables**

**$\Rightarrow$  unique solution.**

## Theorem 2 (Existence and Uniqueness Theorem)

1. A linear system is consistent if and only if the rightmost column of the augmented matrix is not a pivot column, i.e., if and only if an echelon form of the augmented matrix has no row of the form

$$\left[ \begin{array}{cccc} 0 & \dots & 0 & b \end{array} \right] \text{ (where } b \text{ is nonzero).}$$

2. If a linear system is consistent, then the solution contains either

- (i) a unique solution (when there are no free variables) or
- (ii) infinitely many solutions (when there is at least one free variable).

### Using Row Reduction to Solve Linear Systems

1. Write the augmented matrix of the system.
2. Use the row reduction algorithm to obtain an equivalent augmented matrix in echelon form. Decide whether the system is consistent. If not, stop; otherwise go to the next step.
3. Continue row reduction to obtain the reduced echelon form.
4. Write the system of equations corresponding to the matrix obtained in step 3.
5. State the solution by expressing each basic variable in terms of the free variables and declare the free variables.

## EXAMPLE:

a) What is the largest possible number of pivots a  $4 \times 6$  matrix can have? Why?

Four, because there can be at most one pivot position per row.

b) What is the largest possible number of pivots a  $6 \times 4$  matrix can have? Why?

Four, because there can be at most one pivot position per column.

c) How many solutions does a consistent linear system of 3 equations and 4 unknowns have? Why?

An infinite number, because there must be at least one free variable.

d) Suppose the coefficient matrix corresponding to a linear system is  $4 \times 6$  and has 3 pivot columns. How many pivot columns does the augmented matrix have if the linear system is inconsistent?

Four, because:

A system is inconsistent iff the rightmost column of the augmented matrix is a pivot column.