

## 4.3 Linearly Independent Sets; Bases

### Definition

A set of vectors  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$  in a vector space  $V$  is said to be **linearly independent** if the vector equation

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p = \mathbf{0}$$

has only the trivial solution  $c_1 = 0, \dots, c_p = 0$ .

The set  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$  is said to be **linearly dependent** if there exists weights  $c_1, \dots, c_p$ , not all 0, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p = \mathbf{0}.$$

The following results from Section 1.7 are still true for more general vectors spaces.

A set containing the zero vector is linearly dependent.

A set of two vectors is linearly dependent if and only if one is a multiple of the other.

A set containing the zero vector is linearly independent.

**EXAMPLE:**  $\left\{ \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ 3 & 0 \end{bmatrix} \right\}$  is a

linearly \_\_\_\_\_ set.

**EXAMPLE:**  $\left\{ \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, \begin{bmatrix} 3 & 6 \\ 9 & 11 \end{bmatrix} \right\}$  is a linearly

\_\_\_\_\_ set since  $\begin{bmatrix} 3 & 6 \\ 9 & 11 \end{bmatrix}$  is not a

multiple of  $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ .

#### Theorem 4

An indexed set  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$  of two or more vectors, with  $\mathbf{v}_1 \neq \mathbf{0}$ , is linearly dependent if and only if some vector  $\mathbf{v}_j$  ( $j > 1$ ) is a linear combination of the preceding vectors  $\mathbf{v}_1, \dots, \mathbf{v}_{j-1}$ .

**EXAMPLE:** Let  $\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$  be a set of vectors in  $\mathbf{P}_2$  where  $\mathbf{p}_1(t) = t$ ,  $\mathbf{p}_2(t) = t^2$ , and  $\mathbf{p}_3(t) = 4t + 2t^2$ . Is this a linearly dependent set?


*Solution:* Since  $\mathbf{p}_3 = \underline{\hspace{1cm}}\mathbf{p}_1 + \underline{\hspace{1cm}}\mathbf{p}_2$ ,  $\{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$  is a linearly \_\_\_\_\_ set.

## A Basis Set

Let  $H$  be the plane illustrated below. Which of the following are valid descriptions of  $H$ ?

(a)  $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$       (b)  $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_3\}$

(c)  $H = \text{Span}\{\mathbf{v}_2, \mathbf{v}_3\}$       (d)  $H = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$



A *basis set* is an “efficient” spanning set containing no unnecessary vectors. In this case, we would consider the linearly independent sets  $\{\mathbf{v}_1, \mathbf{v}_2\}$  and  $\{\mathbf{v}_1, \mathbf{v}_3\}$  to both be examples of basis sets or bases (plural for basis) for  $H$ .

### DEFINITION

Let  $H$  be a subspace of a vector space  $V$ . An indexed set of vectors  $\beta = \{\mathbf{b}_1, \dots, \mathbf{b}_p\}$  in  $V$  is a basis for  $H$  if

- (i)  $\beta$  is a linearly independent set, and
- (ii)  $H = \text{Span}\{\mathbf{b}_1, \dots, \mathbf{b}_p\}$ .

**EXAMPLE:** Let  $\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ ,  $\mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{e}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ .

Show that  $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  is a basis for  $\mathbf{R}^3$ . The set  $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  is called a **standard basis** for  $\mathbf{R}^3$ .

*Solutions:* (Review the IMT, page 129) Let

$$A = \begin{bmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}. \text{ Since } A \text{ has 3 pivots,}$$

the columns of  $A$  are linearly \_\_\_\_\_ by

the IMT and the columns of  $A$  \_\_\_\_\_

by IMT. Therefore,  $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  is a basis for  $\mathbf{R}^3$ .

**EXAMPLE:** Let  $S = \{1, t, t^2, \dots, t^n\}$ . Show that  $S$  is a basis for  $\mathbf{P}_n$ .

*Solution:* Any polynomial in  $\mathbf{P}_n$  is in span of  $S$ . To show that  $S$  is linearly independent, assume  $c_0 \cdot 1 + c_1 \cdot t + \dots + c_n \cdot t^n = \mathbf{0}$

Then  $c_0 = c_1 = \dots = c_n = 0$ . Hence  $S$  is a basis for  $\mathbf{P}_n$ .

**EXAMPLE:** Let  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix}$ .

Is  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  a **basis** for  $\mathbf{R}^3$ ?

*Solution:* Again, let  $A = [\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3] = \begin{bmatrix} 1 & 0 & 1 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$ . Using row

reduction,

$$\begin{bmatrix} 1 & 0 & 1 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -2 \\ 0 & 1 & 3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 5 \end{bmatrix}$$

and since there are 3 pivots, the columns of  $A$  are linearly independent and they span  $\mathbf{R}^3$  by the IMT. Therefore  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is a **basis** for  $\mathbf{R}^3$ .

**EXAMPLE:** Explain why each of the following sets is **not** a basis for  $\mathbf{R}^3$ .

$$(a) \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 7 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -3 \\ 7 \end{bmatrix} \right\}$$

$$(b) \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \right\}$$

## Bases for Nul A

**EXAMPLE:** Find a basis for Nul A where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

*Solution:* Row reduce  $\begin{bmatrix} A & \mathbf{0} \end{bmatrix}$ :

$$\begin{bmatrix} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{bmatrix}$$

$$x_1 = -2x_2 - 13x_4 - 33x_5$$

$$x_3 = 6x_4 + 15x_5$$

$x_2, x_4$  and  $x_5$  are free

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} -2x_2 - 13x_4 - 33x_5 \\ x_2 \\ 6x_4 + 15x_5 \\ x_4 \\ x_5 \end{bmatrix} =$$
$$x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix}$$

$\uparrow$                        $\uparrow$                        $\uparrow$   
**u**                                      **v**                                      **w**

Therefore  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  is a spanning set for  $\text{Nul } A$ . In the last section we observed that this set is linearly independent. Therefore  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  is a basis for  $\text{Nul } A$ . The technique used here always provides a linearly independent set.

## The Spanning Set Theorem

A basis can be constructed from a spanning set of vectors by discarding vectors which are linear combinations of preceding vectors in the indexed set.

**EXAMPLE:** Suppose  $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 0 \\ -1 \end{bmatrix}$  and

$$\mathbf{v}_3 = \begin{bmatrix} -2 \\ -3 \end{bmatrix}.$$

*Solution:* If  $\mathbf{x}$  is in  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ , then

$$\begin{aligned} \mathbf{x} &= c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3(\text{---}\mathbf{v}_1 + \text{---}\mathbf{v}_2) \\ &= \text{---}\mathbf{v}_1 + \text{---}\mathbf{v}_2 \end{aligned}$$

Therefore,

$$\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\} = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}.$$



## **THEOREM 5    The Spanning Set Theorem**

Let  $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$  be a set in  $V$  and let  $H = \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ .

- a. If one of the vectors in  $S$  - say  $\mathbf{v}_k$  - is a linear combination of the remaining vectors in  $S$ , then the set formed from  $S$  by removing  $\mathbf{v}_k$  still spans  $H$ .
- b. If  $H \neq \{\mathbf{0}\}$ , some subset of  $S$  is a basis for  $H$ .

## Bases for Col A

**EXAMPLE:** Find a basis for Col A, where

$$A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \mathbf{a}_3 \ \mathbf{a}_4] = \begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}.$$

*Solution:* Row reduce:

$$[\mathbf{a}_1 \ \mathbf{a}_2 \ \mathbf{a}_3 \ \mathbf{a}_4] \sim \dots \sim \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & 1 & 5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = [\mathbf{b}_1 \ \mathbf{b}_2 \ \mathbf{b}_3 \ \mathbf{b}_4]$$

Note that

$$\mathbf{b}_2 = \underline{\hspace{1cm}} \mathbf{b}_1 \quad \text{and} \quad \mathbf{a}_2 = \underline{\hspace{1cm}} \mathbf{a}_1$$

$$\mathbf{b}_4 = 4\mathbf{b}_1 + 5\mathbf{b}_3 \quad \text{and} \quad \mathbf{a}_4 = 4\mathbf{a}_1 + 5\mathbf{a}_3$$

$\mathbf{b}_1$  and  $\mathbf{b}_3$  are not multiples of each other

$\mathbf{a}_1$  and  $\mathbf{a}_3$  are not multiples of each other

Elementary row operations on a matrix do not affect the linear dependence relations among the columns of the matrix.

Therefore  $\text{Span}\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3, \mathbf{a}_4\} = \text{Span}\{\mathbf{a}_1, \mathbf{a}_3\}$  and  $\{\mathbf{a}_1, \mathbf{a}_3\}$  is a basis for Col A.

## THEOREM 6

The pivot columns of a matrix  $A$  form a basis for  $\text{Col } A$ .

**EXAMPLE:** Let  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -2 \\ -4 \\ 6 \end{bmatrix}$ ,  $\mathbf{v}_3 = \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}$ .

Find a basis for  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ .

*Solution:* Let  $A = \begin{bmatrix} 1 & -2 & 3 \\ 2 & -4 & 6 \\ -3 & 6 & 9 \end{bmatrix}$  and note that

$$\text{Col } A = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}.$$

By row reduction,  $A \sim \begin{bmatrix} 1 & -2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$ . Therefore a basis

for  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is  $\left\{ \begin{bmatrix} \phantom{1} \\ \phantom{2} \\ \phantom{-3} \end{bmatrix}, \begin{bmatrix} \phantom{-2} \\ \phantom{-4} \\ \phantom{6} \end{bmatrix} \right\}$ .

## Review:

1. To find a basis for  $\text{Nul } A$ , use elementary row operations to transform  $[A \ \mathbf{0}]$  to an equivalent reduced row echelon form  $[B \ \mathbf{0}]$ . Use the reduced row echelon form to find parametric form of the general solution to  $A\mathbf{x} = \mathbf{0}$ . The vectors found in this parametric form of the general solution form a basis for  $\text{Nul } A$ .

2. A basis for  $\text{Col } A$  is formed from the pivot columns of  $A$ .  
**Warning: Use the pivot columns of  $A$ , not the pivot columns of  $B$ , where  $B$  is in reduced echelon form and is row equivalent to  $A$ .**